

Thu1.4

Fincke et al.

## Postponement of saturation of the Richtmyer-Meshkov instability by convergence

**J.R. Fincke<sup>1</sup>, N.E. Lanier<sup>1</sup>, S.H. Batha<sup>1</sup>, J.M. Taccetti<sup>1</sup>, R.M. Hueckstaedt<sup>1</sup>,  
G.R. Magelssen<sup>1</sup>, N.D. Delameter<sup>1</sup>, M.M. Balkey<sup>1</sup>, S.D. Rothman<sup>2</sup>, K.M. Parker<sup>2</sup> & C. J. Horsfield<sup>2</sup>**

1. Los Alamos National Laboratory  
Los Alamos, New Mexico 87545  
[jfincke@lanl.gov](mailto:jfincke@lanl.gov)

2. AWE  
Aldermaston, UK

Strongly driven cylindrically convergent implosions with well characterized surface perturbations were conducted on the OMEGA laser (Broehly, et al (1977)). The cylindrical targets, consisting of a low density foam core and an aluminum shell covered by an epoxy ablator, are directly driven by fifty laser beams ( $18 \pm 0.3$  kJ, 351 nm, 1 ns pulse width). The outer surface of the aluminum shell is machined to form perturbations with wavenumbers ( $k = 2\pi/\lambda$ ,  $\mu\text{m}^{-1}$ )  $0.08 < k < 2.5$  ( $\lambda = 2.5, 9, 25, \text{ and } 75 \mu\text{m}$ ) and initial amplitudes  $0.03 < \eta_0/\lambda < 0.8$ . The perturbations are in the in the r-z plane with r being the radius in cylindrical coordinates and z is the axis of the cylinder. The aluminum shell is calculated to preheat to  $\approx 3$  eV prior to interaction with the Mach 6 shock launched by the laser drive. The Atwood number is  $\approx 0.6$ .

We observe that the perturbations continue to grow approximately linearly, and even exhibit a noticeable increase in growth rate with time well into the amplitude range where saturation is expected in planar geometry. In planar geometry mode saturation and transition to a slow growing spike and bubble configuration has been experimentally observed at  $\eta/\lambda \approx 0.3$  (Dimonte (1993)). We, however, observe no evidence of saturation for an  $\eta/\lambda$  ratio as large as 5. The perturbation growth rate is observed to scale proportionally with k for  $\eta_0 k < 1.4$ , while for  $\eta_0 k \geq 5$  wavenumber scaling is violated and what is likely a transition to turbulent growth is observed. The rate at which the apparent mix width grows, a consequence of both convergence and instability growth, is consistent with Bell's linear theory (Bell (1951)) of perturbation growth in a converging geometry.

### References

- Bell, G. I., 1951 Taylor instability on cylinders and spheres in the small amplitude approximation, LA-1321, LANL.  
Broehly, et al 1977 Initial performance of the OMEGA laser system, Opt. Commun, **133**, 495.  
Dimonte, G. and Remington, B., 1993 Richtmyer-Meshkov Experiments on the Nova Laser at high compression, Phys. Rev. Lett., **70**, 1806.