

Poster 2

Schilling et al.

Investigation of the large-scale and statistical properties of Richtmyer-Meshkov instability-induced mixing

Oleg Schilling¹, Marco Latini², Wai-Sun Don³ & Barna Bihari¹

1. Lawrence Livermore National Laboratory, Livermore, California, USA

schilling1@llnl.gov

2. Applied and Computational Mathematics

California Institute of Technology, Pasadena, California, USA

mlatini@caltech.edu

3. Division of Applied Mathematics

Brown University, Providence, Rhode Island, USA

wsdon@mail.cfm.brown.edu

The weighted essentially non-oscillatory method (Jiang & Shu 1996; Shu 1999) is applied to study the evolution of single- and two-mode Richtmyer-Meshkov instability-induced mixing. The numerical method is validated using two-dimensional simulations of the Mach 1.21 air/SF₆ shock tube experiments of Collins and Jacobs (2002) that include reshock of the evolving interface. The flow structure and perturbation amplitude growth obtained from the simulations are shown to be in very good agreement with the corresponding experimental data. The amplitude growth is also compared to the growth predicted by several nonlinear analytical models.

High-resolution, three-dimensional simulations are performed to study the spatial structure, spectra, and statistics in the Mach 1.5 air/SF₆ shock tube experiment of Vetter and Sturtevant (1995). The interfacial perturbation, including the embedded wire mesh, is modeled following the two-scale representation of Cohen et al. (2002). The mixing layer width is compared to the experimental data. The time-evolution of the kinetic energy spectra corresponding to velocity fluctuations in both the streamwise and spanwise directions and the density variance spectra are investigated. The balance of terms in the kinetic energy and enstrophy evolution equations is studied to provide insight into the baroclinic production and vortex stretching mechanisms that drive the instability and turbulence. Simulations using different orders of spatial accuracy are compared at different grid resolutions to investigate the dependence of quantities on the resolution.

This work was performed under the auspices of the U.S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

References

- Cohen, R. H., Dannevik, W. P., Dimits, A. M., Eliason, D. E., Mirin, A. A., Zhou, Y., Porter, D. H. & Woodward, P. R. 2002 Three-dimensional simulation of a Richtmyer-Meshkov instability with a two-scale initial perturbation; *Phys. Fluids* 14, 3692-3709.
- Collins, B. D. & Jacobs, J. W. 2002 PLIF flow visualization and measurements of the Richtmyer-Meshkov instability of an air/SF₆ interface; *J. Fluid Mech.* 464, 113-136.
- Jiang, G.-S. & Shu, C.-W. 1996 Efficient Implementation of Weighted ENO Schemes; *J. Comp. Phys.* 126, 202-228.
- Shu, C.-W. 1999 High Order ENO and WENO Schemes for Computational Fluid Dynamics; *High-Order Methods for Computational Physics*, edited by T. J. Barth and H. Deconinck, 439-582, Springer-Verlag.
- Vetter, M. & Sturtevant, B. 1995 Experiments on the Richtmyer-Meshkov Instability of an Air/SF₆ Interface; *Shock Waves* 4, 247-252.

UCRL-ABS-201724